

DRUG FACTS

Active ingredient(s)

Purpose

Use(s)

temporarily relieves these cold, sinus and flu symptoms:

- nasal and sinus congestion
- headache
- stuffy nose
- sore throat
- minor aches and pains
- fever

Warnings

Do not use

- in a child under 2 years of age
- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if the child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if you have

- problems or serious side effects from taking pain relievers or fever reducers
- stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain
- ulcers
- bleeding problems
- asthma
- not been drinking fluids
- lost a lot of fluid due to vomiting or diarrhea
- high blood pressure
- heart or kidney disease
- thyroid disease
- diabetes
- taken a diuretic

Ask a doctor or pharmacist before use if you are

- taking any other drug containing a NSAID (prescription or nonprescription)
- taking a blood thinning (anticoagulant) or steroid drug
- under a doctor's care for any serious condition
- taking any other product that contains pseudoephedrine or any other nasal decongestant
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- the child feels faint, vomits blood, or has bloody or black stools. These are signs of stomach bleeding.
- stomach pain or upset gets worse or lasts
- the child does not get any relief within first day (24 hours) of treatment
- symptoms continue or get worse
- fever or pain or nasal congestion gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- the child gets nervous, dizzy, or sleepless or sleepy
- any new symptoms appear

If pregnant or breast-feeding

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Other information

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if the child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing a NSAID [aspirin, ibuprofen, naproxen, or others]
- takes more or for a longer time than directed

Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by a doctor.

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- do not give longer than 10 days, unless directed by a doctor (see WARNINGS)
- **shake well before using**

- find right dose on chart. If possible, use weight to dose; otherwise use age.

- if needed, repeat dose every **6 hours**

- do not use more than **4 times a day**

- replace original bottle cap to maintain child resistance

- measure only with dosing cup provided. Dosing cup to be used with Children's Advil Cold Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

Weight (lb)	Age (yrs)	Dose (teaspoons)
under 24	under 2 yrs	do not use
24-47	2-5 yrs	1 teaspoon
48-95	6-11 yrs	2 teaspoons

- **each teaspoon contains:** sodium 3 mg
- store at room temperature 20-25°C (68-77°F).
- alcohol free
- read all warnings and directions before use. Keep carton.

Inactive ingredients

artificial flavor, carboxymethylcellulose sodium, citric acid monohydrate, edetate disodium, FD&C blue no. 1, FD&C red no. 40, glycerin, microcrystalline cellulose, polysorbate 80, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum